

Product	Label Contents	Dimensions (WxH)
ZONOR (UK)	5L - Booklet	385 x 150 mm

Ringspot (<i>Mycosphaerella brassicicola</i>) (reduction)	Spring or summer applications of 1.0 L/ha ZONOR® made for the control of Light Leaf Spot may also give some reduction in Ringspot.
FIELD BEANS Apply one application of 1.0 L/ha in 200 - 600 Litres water/ha from GS 40 up to 35 days before harvest to control:	
Chocolate spot and bean rust	Apply at first signs of disease from early flowering.
LINSEED Apply one application of 1.0 L/ha in 100 - 400 Litres water/ha after BBCH 20 up to before the capsules turn brown or 35 days before harvest (whichever occurs first) to control:	
Powdery mildew and botrytis (reduction only)	Apply as disease appears usually during warm humid weather from August onwards to achieve control of the mildew infection and a reduction in botrytis

Compatibility

When applying ZONOR® in tank-mixture with other products, follow any instructions as to the order of mixing on any partner product. Where no instructions are given, fully disperse the partner product before adding ZONOR®.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

ZONOR® - MAPP 18650

CONTAINS 250 G/L (25.8 % W/W) TEBUCONAZOLE AS AN OIL IN WATER EMULSION FORMULATION.

ZONOR® IS A BROAD-SPECTRUM SYSTEMIC FUNGICIDE FOR USE IN WINTER & SPRING CROPS OF WHEAT (EXCLUDING DURUM), BARLEY, OATS, RYE, OILSEED RAPE, FIELD BEANS AND LINSEED.

	<p>DANGER Harmful if swallowed. Harmful by inhalation. Causes serious eye damage. Suspected of damaging the unborn child. Very toxic to aquatic life with long-lasting effects. Keep out of reach of children. Wear protective gloves, protective clothing and eye/face protection. IF IN EYES, rinse cautiously with water for several minutes. Remove contact lenses if worn and easy to do so and continue rinsing. Immediately call a POISON CENTRE or doctor/physician. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple-rinsed empty containers which can be disposed of as non-hazardous waste. Do not contaminate water with the product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads). To avoid risks to human health and the environment, comply with the instructions for use.</p>
--	--

IMPORTANT INFORMATION: FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE				
Crops	Maximum individual dose (L/ha)	Maximum No of treatments per crop	Maximum Total dose (L/ha)	Latest timing of application
Wheat, barley oats and rye (winter) after BBCH 30	1.0	2 per crop ¹		End of flowering (BBCH 69)
Oilseed rape: At BBCH 14 – 19 or After BBCH 20	0.5 ⁴		0.5 ²	9 or more leaves unfolded
Field beans after BBCH 40	1.0	1 per crop	1.0 ³	End of flowering
Linseed after BBCH 20	1.0	1 per crop		35 days before harvest
				Before brown capsule stage and 35 days before harvest
Other specific restrictions:				
<ul style="list-style-type: none"> A minimum interval of 14 days must elapse between applications. <ol style="list-style-type: none"> When used on cereals, a maximum total dose of 1.0 L/ha should be applied after BBCH 30 and before early boot stage (BBCH 39). A second application of 1.0 L/ha cannot be applied until after BBCH 40. When used on oilseed rape a maximum total dose of 0.5 L/ha can be applied between growth stages BBCH 14 and BBCH 19. When used on oilseed rape a maximum total dose of 1.0 L/ha can be applied between growth stages BBCH 20 and BBCH 69. When used on oilseed rape, if an application is made between BBCH 14 and BBCH 19 then no further applications are allowed on the crop. Do not apply by hand-held equipment. 				
READ THE LABEL BEFORE USING. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS				

THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK.

APPROVAL HOLDER AND MARKETING COMPANY:

Life Scientific Ltd, Block 4, Belfield Office Park, Beech Hill Road, Dublin 4, Ireland, Tel: +353 1 2832024

FOR 24-HOUR EMERGENCY INFORMATION CONTACT NHS 111.

PROTECT FROM FROST

MADE IN EU

SHAKE WELL BEFORE USE

BATCH NO. SEE PACKAGING



1703202023802_V4

